

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## *Dicamba M 420SL*

Selective Herbicide

### ACTIVE CONSTITUENT:

340 g/L MCPA (present as the dimethylamine salt)  
80 g/L DICAMBA (present as the dimethylamine salt)

GROUP

I

HERBICIDE

**For the control of certain broadleaf weeds in winter cereals, pastures,  
turf and non-crop areas.**

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

Sabakem Pty Ltd ACN 151 682 138  
Level 1, 327-333 Police Road, Mulgrave, VIC 3170  
Tel: 0448 888 960  
[www.sabakem.com](http://www.sabakem.com)

**Contents: 1L, 5L, 20L**

## DIRECTIONS FOR USE

### RESTRAINTS

DO NOT spray when rain seems likely to occur within 4 hours.

DO NOT spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or water logging.

DO NOT spray outside crop growth stages recommended as crop damage may result.

DO NOT apply to crops or grass pastures undersown with clover, lucerne or medics.

DO NOT apply to legume pastures.

DO NOT apply to plants suffering from stress.

CROP & SITUATION	WEEDS	STATE	RATE	CRITICAL COMMENTS
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Z21)	Capeweed, Charlock, Common ice-plant, seedling Docks, Doublegee, (Threecornered Jack, Spiny emex), Hexham scent (Melilotus), Mustards, Soldier thistle, Sorrel seedlings, Tares, Tree hogweed, Turnip weed ( <i>Rapistrum rugosum</i> ), Vetches, Ward's weed, Wild radish, Wild turnip ( <i>Brassica tournefortii</i> ), Wireweed (Hogweed)	All states	1L/ha	Spray when most weeds are in the 2-4 leaf stage. In some instances suppression only of wireweed (hogweed), capeweed and wild radish will occur.
	Seedling Cotton thistle, Fat hen	Tas only		
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Z21) until the fully tillered stage and before jointing occurs (Z30)	Cotton thistle to bud formation	Tas only	1.7L/ha	Spray when most weeds are still in the rosette or seedling stage. N.B. Wild radish may need to be resprayed (with 2,4-D or MCPA) if future germinations occur after spraying. In some instances, suppression only of Wireweed (hogweed), Hoary cress and Climbing buckwheat will be obtained.
	Mintweed, New Zealand spinach	Qld, NSW only		
	Capeweed, Charlock, Chickweed, Climbing buckwheat (Black bindweed), Clover, Common ice-plant, seedling Docks, Doublegee (Three cornered Jack, Spiny emex), Fat hen, Hexham scent (Melilotus), seedling Hoary cress, Mouse-eared Chickweed, Mustards, Narrow-leaved plaintain, Saffron thistle, Spurry, Swinecress, Tares, Tree hogweed, Turnip weed ( <i>Rapistrum rugosum</i> ), Variegated thistle, Vetch, Wild radish, Wild turnip ( <i>Brassica tournefortii</i> ), Wireweed (Hogweed)	Qld, NSW Tas, WA, Vic only	1.4L/ha	
Barley, Commercial Rye, Triticale From early tillering stage (Z21) until the fully tillered stage before jointing occurs (Z30)	Volunteer Peas	SA only	1L/ha plus 60mL/ha Dicamba 500g/L	Apply when most peas are 5-7 cm high.
			1L/ha plus 120mL/ha Dicamba 500g/L	

CROP & SITUATION	WEEDS	STATE	RATE			CRITICAL COMMENTS
			Per Hectare	Per 100L	Per 15L Knapsack	
Grass Pastures	Bathurst burr, Caltrop (Yellow vine), Chicory, Crofton weed, Cudweed, Dandelion, Deadnettle, Fleabanes, Fumitory, Groundsel, Heliotrope, Inkweed, Mintweed, Nettles, Noogoora burr, Pennycress, Potato weed, Ragwort, Ripplewort, Shepherd's purse, Smartweed, Soft rush, Stinkwort, Thistles including Cockspur, Scotch (Spear), Slender, Star, Variegated, Wireweed (Hogweed)	All states	2.8-4L	190-270mL	60mL	Use the higher rate on larger weeds.
	Cotton thistle	Tas only				
	Buttercups, Hoary cress seedlings, Horehound, Khaki weed, Purple-top, Veined verbena	All states	4-7L	470mL	100mL	
Turf	Bindy-eye, Capeweed, Catsear, Chickweed, Clovers, Cotula, Creeping oxalis, Cudweeds, Dandelion, Dock, Fleabanes, Fumitory, Jo-Jo (Onehunga), Lamb's tongue, Medics, Mustards, Narrow-leaved plaintain, Pearlwort, Pennywort, Peppergrass, Radish, Scarlet pimpernel, Slender thistle, Soldier thistle, Spear thistle, Toad rush, Variegated thistle, Wild turnip, Wireweed (Hogweed).	All states	<b>Golf Tee, Golf Green:</b> 100mL in 15L water per 150m <sup>2</sup> <b>Bowling Green:</b> 900mL in 200L water per 1400m <sup>2</sup> <b>Large Turf Area:</b> (Field, Park, Golf Fairway) 6.5L in 250-400L water per 10,000m <sup>2</sup> (1 hectare)			Apply to actively growing weeds and moist soil. DO NOT use on Lippia, Strawberry clover or Buffalo grass. DO NOT mow turf for two days before or after application or fertilize within two weeks of spraying.
	Fat hen, Purple-top, Red flowered mallow, Staggerweed, Swinecress	Qld, NSW, Tas, SA, WA only				
Non-Crop Situations	Artichoke, Bindweed, Fennel, Hemlock, Thistle seedlings	All states	5.2L	350mL	80mL	A repeat spray may be necessary.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD:**

**DO NOT GRAZE OR CUT FOR STOCK FOOD 7 DAYS AFTER APPLICATION.**

**HARVEST WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS

GROUP	I	HERBICIDE
-------	---	-----------

## RESISTANT WEEDS WARNING

SABAKEM® DICAMBA M 420SL SELECTIVE HERBICIDE contains members of the Benzoic acid and Phenoxy groups of herbicides. SABAKEM® DICAMBA M 420SL SELECTIVE HERBICIDE has the Disruptors of plant cell growth mode of action. For weed resistance management SABAKEM® DICAMBA M 420SL SELECTIVE HERBICIDE is a Group I Herbicide. Some naturally-occurring weed biotypes resistant to SABAKEM® DICAMBA M 420SL SELECTIVE HERBICIDE and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by SABAKEM® DICAMBA M 420SL SELECTIVE HERBICIDE or any other Group I Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, SABAKEM Pty Ltd accepts no liability for any losses that may result from the failure of SABAKEM® DICAMBA M 420SL SELECTIVE HERBICIDE to control resistant weeds.

## MIXING

Add the required amount of SABAKEM® DICAMBA M 420SL SELECTIVE HERBICIDE directly to the tank and mix well.

## APPLICATION

**For Boom Application:** Apply by a properly calibrated boom spray using not less than 50L of water per hectare unless indicated otherwise in the Critical Comments column.

**For Aerial Application:** Apply at least 25L total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.

**For High Volume Application:** Apply minimum 1500L per hectare.

**For Knapsack Application:** A 15L knapsack treats 150 square metres.

**NOTE:** Spray when weeds are young and actively growing. A temporary wilting may be evident in crops after application. If seasonal conditions are unusually late or dry seek further advice as damage may result from spraying under these conditions. Growers should seek advice before spraying recently released cereal varieties. There is a nil tolerance of pea seed contaminates in cereal grain and less than 100% control of Volunteer Peas could result in dockage.

## COMPATIBILITY

SABAKEM® DICAMBA M 420SL SELECTIVE HERBICIDE is compatible with Flowable Igran<sup>†</sup> (Terbutryn 500SC).

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Spray equipment must be calibrated accurately prior to working on crops. Minimise spray drift by using low pressure and nozzles which do not give a fine droplet size. DO NOT use in high wind. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals. DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots. Keep container sealed (airtight) when not in use. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticulture crops.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, triple rinse, break, crush or puncture and deliver empty packaging for appropriate disposal at an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product must not be burnt.

## SAFETY DIRECTIONS

Harmful if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes and skin. DO NOT inhale vapour. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves, face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with water. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26.

## MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet.

## CONDITIONS OF SALE

The use of this product is beyond the control of Sabakem Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Sabakem Pty Ltd accepts no responsibility for any consequences

resulting from the use of this product. SBAKEM Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

**For specialist advice in an emergency dial  
1 800 638 556  
24 hours Australia wide.**

APVMA Approval No: 69159/ 60096

<sup>†</sup> Not Registered trade mark of Sabakem Pty Ltd

© Sabakem is the registered trade mark of Sabakem Pty Ltd