

Captan 800 WG[®]

Sabakem

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Sabakem

Captan 800 WG[®]

Fungicide

ACTIVE CONSTITUENT: 800 g/kg CAPTAN

GROUP **M4** FUNGICIDE

For the control of certain diseases in a range of fruit crops, turf and ornamentals as shown under Directions for Use.


IMPORTANT: READ THE FULL DIRECTIONS BEFORE OPENING OR USING THIS PRODUCT

NET CONTENTS: 10KG



5H4/Z14/S12
CN/330351
PI:015

UN No. 3077	
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S (Contains CAPTAN)	
PG III	HAZCHEM 2<Z>
In a Transport Emergency DIAL 000 Police or Fire Brigade	



Sabakem Pty Ltd ABN 34 151 682 138
Suite 809, Level 8, 2 Queen Street, Melbourne VIC 3000
Tel: 03 9629 3979 • Fax: 1300 242 436
www.sabakem.com



DOM:

Batch



DIRECTIONS FOR USE					
TREE AND VINE CROPS					
RATE					CRITICAL COMMENTS
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/ Application section.					
CROP	DISEASE	STATE	RATE/ 100L	WHP	
Apples	Black Spot (<i>Venturia inaequalis</i>)	All States	Dilute spraying: 125 g/100L Concentrate spraying: Refer to the Mixing/ Application section	7 days	Apply from pink stage onwards following a copper spray at green tip, then at 7 day intervals until petal fall. After petal fall apply as a cover spray at 10-14 day intervals. The risk of russet may be increased in some varieties. Where this is of concern use alternative materials between full bloom and second cover. Apply to point of run-off. Apply no more than five applications per season. Apply at 50% bud-burst and repeat application 10-14 days later. Apply to point of run-off. Apply no more than five applications per season. Apply in a program of sprays applied at early flowering, 80-100% cap fall, pre-bunch closure, veraison and pre-harvest. Apply to point of run-off. Apply no more than five applications per season. Apply from flowering onwards, at intervals of 10-14 days. Apply to point of run-off. Apply no more than five applications per season. Commence application at early budburst and repeat at 7 to 14 day intervals until prior to flowering to maintain a cover of new grape foliage. Apply no more than 5 applications per season. Use the shorter application interval where there is rapid shoot growth and conditions are favourable for Phomopsis infection (wet weather). Adjust spray application equipment to obtain a thorough coverage of grape buds and shoots.
Pears	Black Spot (<i>Venturia pirina</i>)				
Grapes	Black Spot (<i>Eisinoe ampelina</i>)				
	Grey Mould (<i>Botrytis cinerea</i>)				
	Downy Mildew (<i>Plasmopara viticola</i>)				
	Phomopsis Cane and Leaf Blight (<i>Phomopsis viticola</i>)				
Stone Fruit (except Apricots)	Blossom Blight and Brown Rot (<i>Sclerotinia laxa</i> , <i>S. fructicola</i>)				Apply at any of the following times as part of a spray program to control blossom blight and brown rot – pink bud, 10% blossom, full bloom, petal fall and shuck fall and pre-harvest sprays at 6, 3 and 1 week prior to harvest. Apply to point of run-off. Apply no more than five applications per season. DO NOT apply to apricots as damage to fruit and foliage may occur.
NON TREE AND VINE CROPS					
CROP	DISEASE	STATE	RATE 100/L	WHP	CRITICAL COMMENTS
Strawberries	Grey Mould (<i>Botrytis cinerea</i>), Gloeosporium Fruit Rot	All States	125 g	1 day	Apply every 10 days commencing at blossom stage. Apply to point of run-off. Apply no more than five applications per season.
	Phytophthora Fruit Rot (<i>Phytophthora nicotianae</i> var <i>parasitica</i>), Black Spot (Anthracnose) (<i>Colletotrichum acutatum</i>), Scorch (<i>Diplocarpon earlianum</i>), Leaf Blight (<i>Dendrophoma obscurans</i>)		200 g		
Turf	Brown Patch (<i>Rhizoctonia solani</i>)		25-50 L water/100-150 g in 100 m ² turf	–	For preventative treatment from September to April, apply every 5-6 weeks. Apply no more than five applications per season.
Seedlings (Ornamentals and Tobacco)	Damping-off (<i>Pythium</i> spp.)		125 g		Use 3L of solution per m ² of seedbed. Apply every 7-14 days when weather conditions favouring the disease persists, paying particular attention to bases of seedlings. When treating tobacco seedbeds, wait until seedlings are at least 5cm high.
Roses	Black Spot (<i>Diplocarpon rosae</i>)				Apply to thoroughly wet both surfaces of foliage. Apply every 7-14 days. Use the shorter interval during showery weather. Apply no more than five applications per season.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THE LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

STRAWBERRIES: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.
APPLES, PEARS, GRAPES, STONE FRUIT: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.
DO NOT USE TREATED PRODUCE FOR STOCK FOOD WITHIN 7 DAYS OF TREATMENT.

GENERAL INSTRUCTIONS

FUNGICIDE RESISTANCE WARNING

For fungicide resistance management Sabakem Captan 800 WG[®] Fungicide is a Group M4 Fungicide. Some naturally occurring individual fungi resistant to Sabakem Captan 800 WG[®] Fungicide and other Group M4 Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Sabakem Captan 800 WG[®] Fungicide or other Group M4 Fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Sabakem Pty Ltd accepts no liability for any losses that may result from the failure of Sabakem Sabakem Captan 800 WG[®] Fungicide to control resistant fungi.

GROUP M4 FUNGICIDE

MIXING

Sabakem Captan 800 WG[®] Fungicide mixes readily with water and should not be pre-mixed. Slowly add the required amount of product to the water under agitation in the spray vat. Maintain agitation throughout spraying. After a stoppage, thoroughly re-mix before recommencing spraying.

COMPATIBILITY AND CROP SAFETY

Mixtures of Sabakem Captan 800 WG[®] Fungicide with commonly used wettable powders and suspension concentrate insecticides and fungicides should not cause any problems. Apart from the products listed below mixtures with emulsifiable concentrates are not recommended. If tank mixes are to be used observe all directions, precautions and limitations on all products to be used. Caution should also be exercised when spraying any mixtures on varieties of fruit trees or vines where there has been no previous experience of mixtures with Sabakem Captan 800 WG[®] Fungicide. Sabakem Captan 800 WG[®] Fungicide is compatible with any one of the following products: Anvil[®], Bayfidant[®], Topaz[®], Pennicap-M[®], Gusathion[®] 200SC with or without calcium nitrate, Bugmaster[®] Flowable, Confidor[®], Hymal, Insegar[®], Kumulus[®], Lebaycid[®], Lorsban[®], Mavrik[®] Aquaflo, Omite[®] 300WG (apples only), Pirimor[®] WG, Pyranica[®], Saboteur[®], Sanmitel[®], Systhane[®]. A mixture of Sabakem Captan 800 WG[®] Fungicide with more than one of these products or with any other product may be ineffective or may cause serious damage. The use of such a mixture is not recommended without prior advice, and would therefore be entirely at the user's risk. If other mixtures or 3 way mixtures of any of these products are being considered consult your Sabakem Pty Ltd dealer or representative for advice. Mixtures of A mixture of Sabakem Captan 800 WG[®] Fungicide with Lorsban[®] 500EC or Nimrod[®] may leave an oily residue on the inside of spray vats which may be difficult to remove. A mixture of Sabakem Captan 800 WG[®] Fungicide should not be combined with Bordeaux mixture, spray lime, lime sulphur or oils. DO NOT apply A mixture of Sabakem Captan 800 WG[®] Fungicide in combination with or within 14 days before or after winter or summer oil sprays. Combinations of a mixture of Sabakem Captan 800 WG[®] Fungicide and sulphur should not be used on crops sensitive to sulphur. Sabakem Captan 800 WG[®] Fungicide mixed with Folidol[®] M500 may be phytotoxic to many varieties of apples and pears. If it is necessary to use both Sabakem Captan 800 WG[®] Fungicide and Folidol[®] M500 in a spray program, sprays of these products must be at least 7 days apart. On pears tank mixtures of Sabakem Captan 800 WG[®] Fungicide plus Omite[®] 300 WG and Sabakem Captan 800 WG[®] Fungicide and Gusathion[®] (without calcium nitrate) plus Omite[®] 300WG have been found to be phytotoxic. DO NOT tank mix these products, for use on pears. Caution should be taken when applying chemical mixtures to all crops. Varieties within crop groups may react differently therefore a small area should be tested prior to application to crop.

APPLICATION

Ground application: These rates are for high volume spraying to point of run-off. DO NOT apply by aircraft.

Dilute Spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way.

Example Only:

- Dilute spray volume as determined above: For example 1000 L/ha
 - Your chosen concentrate spray volume: For example 500 L/ha
 - The concentration factor in this example is: 2 x (i.e. 1000 L ÷ 500L = 2)
 - If the dilute label rate is 125 g/100 L, then the concentrate rate becomes 2 x 125, that is 250 g/100 L of concentrate spray.
- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
 - DO NOT use at a concentration rate higher than that specified in the Critical Comments as crop safety has not been assessed.
 - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

PRECAUTION

RE-ENTRY PERIODS: DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT apply directly to water. DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto water, or to areas where surface water is present. DO NOT contaminate surface water with the concentrated product, spray mix, rinsate, drainage water from glasshouses or the used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Avoid preparation of excess spray, but if necessary, spray out excess mixed spray or dispose of in a soil pit which does not drain into waterways or ponds. Degradation of excess material may be hastened by making solutions alkaline with lime or washing soda. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will damage eyes. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. Repeated exposure may cause allergic disorders. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26.

SAFETY DATA SHEET

If additional hazard information is required refer to the Safety Data Sheet.

CONDITIONS OF SALE: The use of this product is beyond the control of Sabakem Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Sabakem Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Sabakem Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: May cause an allergic skin reaction. Causes serious eye damage. Toxic if inhaled. Suspected of causing cancer. Very toxic to aquatic life.

© Registered trademark of Sabakem Pty Ltd

† Not a trademark of Sabakem Pty Ltd

APVMA Approval No.: 68909 / 59362

Sabakem Pty Ltd ABN 34 151 682 138
 Suite 809, Level 8, 2 Queen Street, Melbourne VIC 3000
 Tel: 03 9629 3979 • Fax: 1300 242 436
 • www.sabakem.com