

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Sabakem

Chlorothalonil 720 SC®

Fungicide

ACTIVE CONSTITUENT: 720 g/L CHLOROTHALONIL

GROUP **M5** FUNGICIDE

For the control of fungal diseases on almonds, bananas, cucurbits, grapes, ornamentals, peanuts, stone fruit, tobacco and vegetables as specified in the Directions for Use.

IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING THIS PRODUCT

Sabakem Pty Ltd ABN 34 151 682 138
Suite 809, Level 8, 2 Queen Street, Melbourne VIC 3000
Tel: 03 9629 3979 • Fax: 1300 242 436
www.sabakem.com

Leaflet_0120

APVMA Approval No: 86767 / 116529

GENERAL INSTRUCTIONS

Sabakem Chlorothalonil 720 SC Fungicide is a protectant fungicide only, it must be applied before disease occurs/becomes established to be highly effective. Use disease warning services, or past history to determine timing of the first spray. If diseases are already present, then an eradicant product should be used first and mixed with Sabakem Chlorothalonil 720 SC Fungicide.

MIXING

Invert the container several times before opening. Half fill spray vat and with agitation system running, add the required amount of product.

APPLICATION

Dilute Spraying

- Use sprayer designed to apply high volume of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines and expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- Use sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way.

Example Only

- Dilute spray volume as determined above: for example 1,500 L/ha.
 - Your chosen concentrate spray volume: for example 500 L/ha.
 - The concentration factor in this example is: 3X (i.e. 1,500 L / 500 L = 3)
 - If the dilute label rate is 40 mL/100 L, then the concentrate rate becomes 3 x 40, that is 120 mL/100L of concentrate spray.
- The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to change as the crop grows.
 - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow Best Practice.

COMPATIBILITY

This product is compatible with wettable powder formulations of the most commonly used fungicides, insecticides and miticides.

DO NOT combine with oil-based emulsifiable or flowable pesticides, unless prior experience has shown the combination to be non-injurious to your crop species.

This product should not be mixed with spraying oils or sprayed onto crops that have been sprayed with oil for at least 10 days after the oil spray. Oils should not be sprayed on crops treated with this product for at least 10 days after the last spray. Wetting agents have does not improved performance. Under some conditions, certain surfactant can cause plant injury.

FUNGICIDE RESISTANCE WARNING

Sabakem Chlorothalonil 720 SC Fungicide is a member of the multi-site activity group of fungicides. For fungicide resistance management Sabakem Chlorothalonil 720 SC Fungicide is a Group M5 Fungicide. Some naturally-occurring individual fungi resistant to Sabakem Chlorothalonil 720 SC Fungicide and other Group M5 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. The resistant fungi will not be controlled by Sabakem Chlorothalonil 720 SC Fungicide or other Group M5 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Sabakem Pty Ltd accepts no liability for any losses that may result from failure of Sabakem Chlorothalonil 720 SC Fungicide to control resistant fungi.

PROTECTION OF LIVESTOCK

DO NOT contaminate feed or foodstuffs with this product or empty containers.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Attacks eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. Repeated exposure may cause allergic disorders. Sensitive workers should wear protective clothing. When preparing spray and using prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and goggles and disposable mist mask. If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product on skin immediately wash area with soap and water. If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from the supplier.

CONDITIONS OF SALE

The use of Sabakem Chlorothalonil 720 SC Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Sabakem Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Sabakem Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia:
Causes serious eye damage. Fatal if inhaled. Suspected of causing cancer.
Very toxic to aquatic life with long lasting effects.

© Registered trademark of Sabakem Pty Ltd

DIRECTIONS FOR USE

RESTRAINTS: DO NOT mix spraying oils or wetting agents with this product.

For all stone fruit except Cherries, the following additional restraints apply:

DO NOT use oils after bud-swell and during the season if Sabakem Chlorothalonil 720 SC Fungicide is to be used after shuck fall.

DO NOT tank mix this product with EC formulations when spraying after shuck fall.

FRUIT AND VEGETABLES

SITUATION & CROP	DISEASE	STATE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Artichoke-Globe	Grey Mould (<i>Botrytis cinerea</i>)	All States	1.8-2.3 L/ha	1	Apply at 7 to 10 day intervals as required. Use the higher rate if disease pressure is high.
Bananas DO NOT apply to bananas within 14 days of applying an oil based spray or crop damage will occur	Leaf Spot (<i>Mycosphaerella musicola</i>)	Qld, NSW, NT only	1.8 L/ha	1	North Qld: Spray every 14 days late January to end of March. Extend to 21 days after March and then 28-35 days in the drier months of the year. South Qld: Spray every 21 days from start of December to end of April. WA: Spray every 10 to 14 days from November to end of May and extend to 14 - 21 days from May to November. NSW: Spray every 21 days from end of November to end of May.
		WA only	1.5 L/ha		
Broccoli, Brussels sprouts,	Ring Spot (<i>Mycosphaerella brassicicola</i>)	All States	1.8-3.5 L/ha	7	Make the first application as soon as conditions favour development of the disease. Repeat at 14 day intervals. Use higher rate if disease pressure is high.
			1.8-2.3 L/ha	1	
Capsicums Peppers	Grey Mould (<i>Botrytis cinerea</i>), Anthracnose (<i>Colletotrichum acutatum</i>)		1.8-2.3 L/ha	1	Apply at 7 to 10 day intervals as required. Use the higher rate if disease pressure is high.
Cabbages, Cauliflowers	Ring Spot (<i>Mycosphaerella brassicicola</i>)		1.8-3.5 L/ha	7	Make the first application a soon as conditions favour disease development of the disease. Repeat at 14 day intervals. Use the higher rate if disease pressure is high.
Carrots	Leaf Blight (<i>Alternaria dauci</i>)		1.8 L/ha	7	Commence application as soon as symptoms favour the development of the disease and then spray every 7-14 days.
Celery	Septoria Leaf Spot (<i>Septoria apiicola</i>)	All States	1.8 L/ha	1	Apply every 7-14 days. Apply also to seedbeds.
	Cercospora Early Blight (<i>Cercospora apii</i>)	NSW, WA only	1.2 L/ha		Apply every 10 to 14 days.
Cucurbits including Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, Rockmelon, Watermelon, Zucchini	Downy Mildew (<i>Pseudoperonospora cubensis</i>), Gummy Stem Blight (<i>Didymella bryoniae</i>), Anthracnose (<i>Colletotrichum orbiculare</i>), Leaf Blight (<i>Alternaria cucumerina</i>), Target Leaf Spot (<i>Cercospora citrullina</i>)	All States	1.6-2.5 L/ha	1	Apply on a 7 to 14 day schedule. Apply rates towards the higher end of the recommended range at 7 day intervals when disease pressure is high. Commence spray program early at first true leaf and before vining or flowering. When used at these rates Sabakem Chlorothalonil 720 SC Fungicide will suppress Powdery Mildew.
	Belly Rot (<i>Rhizoctonia solani</i>)				If applying Sabakem Chlorothalonil 720 SC Fungicide for the above foliar diseases, useful suppression of Belly Rot will be obtained where fruit form on soil covered by earlier Sabakem Chlorothalonil 720 SC Fungicide spraying.
Endive	Grey Mould (<i>Botrytis cinerea</i>)		1.8-2.3 L/ha		Apply at 7 to 10 day intervals as required. Use the higher rate if disease pressure is high.
Leeks Shallots	Purple Blotch (<i>Alternaria porri</i>)				Make the first application as soon as conditions favour development of the disease. Repeat at 7 to 10 day intervals, particularly during wet weather.
Okra	Leaf Disease/Spot	Qld, WA only			Make the first application as soon as conditions favour development of the disease. Repeat at 7 to 10 day intervals.
Onions (excluding spring onions)	Downy Mildew (<i>Peronospora destructor</i>)	All States	2.3 L/ha	14	Make the first application as soon as conditions favour development of the disease. Repeat at 14 day intervals.
Potato	Early Blight or Target Spot (<i>Alternaria solani</i>), Late Blight or Irish Blight (<i>Phytophthora infestans</i>)		1.1-1.8 L/ha	-	Commence application at flowering time or earlier if weather conditions are favourable for disease infection. Spray every 7 to 14 days until within 14 days of desiccation or harvest.
Rhubarb	Ascochyta Leaf Spot (<i>Ascochyta rhei</i>)		140 mL/100 L	7	Apply every 10 to 14 days.
Radish	Grey Mould (<i>Botrytis cinerea</i>)	All States	1.8-2.3 L/ha	1	Apply every 7 to 10 day intervals as required. Use the higher rate if disease pressure is high.
Sweet Corn	Turcicum Leaf Blight (<i>Exserohilum turcicum</i>)				Make the first application as soon as conditions favour development of the disease. Repeat at 7 to 10 day intervals.
Tomato	Early Blight or Target Spot (<i>Alternaria solani</i>), Leaf Mould (<i>Fulvia fulva</i>), Grey Leaf Spot (<i>Stemphylium solani</i>), Botrytis Rot or Grey Mould (<i>Botrytis cinerea</i>)				Make the first application as soon as conditions favour development of the disease. Spray at 7 to 14 day intervals. Continue into the picking season to obtain best results against Leaf Mould and Botrytis Rot.
		Qld, WA only			Apply every 7 to 10 day intervals as required. Use the higher rate if disease pressure is high.

LEGUMES

SITUATION & CROP	DISEASE	STATE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Beans – Broad (Faba)	Chocolate Spot (<i>Botrytis fabae</i>), Rust (<i>Uromyces viciae-fabae</i>)	All States	1.4-2.3 L/ha	7	Make the first application as soon as conditions favour development of the disease. Repeat at 7 to 14 day intervals. Use the higher rate if the crop is dense and disease pressure is high.
Peanuts	Early Leaf Spot (<i>Cercospora arachidicola</i>), Late Leaf Spot (<i>Cercosporidium personatum</i>), Peppery Leaf Spot (<i>Leptosphaerulina trifolii</i>), Peanut Rust (<i>Puccinia arachidis</i>), Net Blotch (<i>Didymosphaeria arachidicola</i>)	Qld, NSW, WA, NT only	1.1-1.8 L/ha	-	Use higher rate for peanut rust. South Burnett, NSW, WA: Commence application as soon as conditions favour development of the disease, repeat as necessary to maintain control. Atherton Tablelands: Commence application 6 weeks after planting and repeat every 2 weeks.

LEGUMES - continued

SITUATION & CROP	DISEASE	STATE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Peas	Downy Mildew (<i>Peronospora viciae</i>)	Qld, Tas, WA only	1.1-1.8 L/ha	7	Make the first application as soon as conditions favour development of the disease. Repeat at 7 to 14 day intervals. Use the higher rate if the crop is dense and disease pressure is high.

MISCELLANEOUS					
SITUATION & CROP	DISEASE	STATE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Ornamentals Chrysanthemums, Geraniums, Iris & other ornamentals	Botrytis, Grey Mould, Rust diseases, Septoria and Alternaria Leaf Spot diseases	All States	100 mL/100 L	-	Apply at 7-14 day intervals. Use 7 day interval under severe conditions. Avoid application to blooms in Azalea, Hydrangea and Petunia.
Tobacco	Tobacco Leaf Spot (<i>Ascochyta arida</i>)	Qld, WA only	220 mL/100 L	-	Apply every 7 days in seedbed application only.

TREE AND VINE CROPS					
RATE					CRITICAL COMMENTS
This table shows rates for dilute spraying. For concentrate spraying refer to the Mixing/Application Section.					For uses in the table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Ensure thorough coverage.
SITUATION & CROP	DISEASE	STATE	RATE	WHP (DAYS)	
Almonds	Shot-hole (<i>Stigmina carpophila</i>), Stone Fruit Rust (<i>Tranzschelia discolor</i>)	NSW, Vic, Tas, SA, WA only	160 mL/100 L	-	Apply at bud-swell, bud-burst, pink-bud, shuck fall, cap fall. Apply every 10 to 14 days. Apply 1 week pre-harvest.
Apricots	Brown Rot – Fruit (<i>Monilinia fructicola</i>), Blossom Blight (<i>Monilinia laxa</i>)			7	Apply at bud-swell, bud-burst, pink bud, full bloom. Applications later than 35 days prior to harvest may result in unacceptable fruit skin damage. See also Additional Restraints for stone fruit.
	Shot-hole (<i>Stigmina carpophila</i>), Stone Fruit Rust (<i>Tranzschelia discolor</i>), Freckle (<i>Venturia carpophila</i>)				Apply at bud-swell, bud-burst, pink bud, shuck fall, cap fall, then every 10 to 14 days.
Cherries	Brown Rot – Fruit (<i>Monilinia fructicola</i>), Blossom Blight (<i>Monilinia laxa</i>)	All States			Apply at bud-swell, bud burst, pink bud, early blossom, and full bloom. Apply 3 weeks pre-harvest and 1 week pre-harvest.
	Shot-hole (<i>Stigmina carpophila</i>), Stone Fruit Rust (<i>Tranzschelia discolor</i>)				Apply at bud-swell, bud burst, pink bud, shuck fall, cap fall, then every 10 to 14 days. Apply 1 week pre-harvest.
	Transit Rot (<i>Rhizopus stolonifer</i>)				Apply 1 week pre-harvest.
Grapes Note: russetting of some table grape varieties may occur	Downy Mildew (<i>Plasmopara viticola</i>), Bunch Rot (<i>Botrytis cinerea</i>)		160-210 mL/100 L	Dessert 7 Wine 14	Spray at first appearance of the foliage disease. Apply at 7-14 day intervals particularly during wet weather and periods of rapid growth. For Bunch Rot, applications at early full bloom, two weeks later, at veraison and two weeks prior to harvest. Table grapes: russetting of some varieties may occur. DO NOT apply after cap fall on these varieties.
	Black Spot (Anthracnose) (<i>Elsinoe ampelina</i>)		210 mL/100 L		If Black Spot has occurred in the vineyard in any of the last three years, apply at budburst and 14 days later when shoots are 10-15 cm long. If wet/humid conditions persist spray again when shoots are 20-30 cm long to protect new growth. Some initial minimal leaf damage on the new shoots of sensitive varieties can occur if spray treatments are applied at budburst and in the first 14 days after budburst. In vineyards with no history of Black Spot, make the first application as soon as conditions favour the development of the disease on the foliage and then at 7-14 day intervals if wet/humid conditions persist. Applications up to cap-fall will not cause russetting in table grape and dried fruit varieties
Nectarines	Shot-hole (<i>Stigmina carpophila</i>)	NSW, Vic, Tas, SA, WA only	160 mL/100 L	7	Apply at bud-swell, pink bud, shuck fall and then repeat each 14 days if weather conditions favour disease. Applications later than 35 days prior to harvest may result in unacceptable fruit skin damage. See also Additional Restraints for stone fruits.
	Brown Rot – Fruit (<i>Monilinia fructicola</i>), Blossom Blight (<i>Monilinia laxa</i>)				Apply at bud-swell, bud burst, pink bud, early blossom and full bloom.
Peaches	Brown Rot – Fruit (<i>Monilinia fructicola</i>), Blossom Blight (<i>Monilinia laxa</i>)	All States	160 mL/100 L	7	Apply at bud-swell, pink bud, early blossom and full bloom. Applications later than 35 days prior to harvest may result in unacceptable fruit skin damage. See also Additional Restraints for stone fruits.
	Shot-hole (<i>Stigmina carpophila</i>), Stone Fruit Rust (<i>Tranzschelia discolor</i>)				Apply at bud-swell and continue at 7-14 day intervals.
	Leaf Curl (<i>Taphrina deformans</i>)				Apply at bud-swell and continue at 7-14 day intervals. In Qld spray only at bud-swell.
Plums	Brown Rot – Fruit (<i>Monilinia fructicola</i>), Blossom Blight (<i>Monilinia laxa</i>) (NOT WA)			1	Apply at bud-swell, bud-burst, early blossom, pink bud and full bloom.
	Shot-hole (<i>Stigmina carpophila</i>), Stone Fruit Rust (<i>Tranzschelia discolor</i>)				Apply at bud-swell, bud burst, shuck fall and cap fall.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Artichokes, Bananas, Capsicums, Celery, Cucurbits, Endive, Leek, Okra, Peppers, Plums, Radish, Shallots, Sweet Corn, Tomatoes, Watercress: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

Apricots, Beans, Broccoli, Brussels Sprouts, Cabbages, Carrots, Cauliflowers, Cherries, Nectarines, Peaches, Peas, Rhubarb, Table Grapes: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

Onions, Wine Grapes: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

Beans, Peanuts, Peas: DO NOT GRAZE LIVESTOCK ON TREATED CROPS.

Almonds, Potatoes, Peanuts (Harvest): NOT REQUIRED WHEN USED AS DIRECTED.