Product Name: Sabakem Glufosinate 400 Herbicide

APVMA Approval No: 88062/132432

Other Limitations:



Label Name:	Sabakem Glufosinate 400 Herbicide
Signal Headings:	CAUTION
	KEEP OUT OF REACH OF CHILDREN
	READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	400 g/L GLUFOSINATE-AMMONIUM
Mode of Action:	GROUP 10 HERBICIDE
Statement of Claims:	For the Non-Residual control of Broadleaf and Grass Weeds in Various Situations as indicated in the Directions for Use
Net Contents:	1L - 1000 L
Restraints:	DO NOT apply by aircraft.
	DO NOT apply when rain is expected within 6 hours. DO NOT apply to weeds under stress due to, for example, very dry, very wet, frosty or
	diseased conditions.
	DO NOT apply under hot dry conditions (temperatures above 33°C with a relative humidity below 50%).
	DO NOT apply greater than 30 L of spray mixture per hectare through OSST equipment (1.44 L of product per hectare).
Directions for Use:	This section contains file attachment.

Withholding Periods:

HARVEST (H)

Avocado, banana, feijoa, guava, kiwifruit, litchi, mango, pawpaw, passionfruit, pineapple, rambutan, blackberry, boysenberry, loganberry, raspberry, citrus fruit, grapes, strawberries, tomatoes, tree nuts: NOT REQUIRED WHEN USED AS DIRECTED.

Pome and stone fruit, olives: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

GRAZING (G)

DO NOT GRAZE OR CUT TREATED AREAS FOR STOCKFOOD FOR 8 WEEKS AFTER APPLICATION.

Trade Advice:

Export of Treated Produce

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with Sabakem Glufosinate 400 Herbicide. If you are growing produce for export, please check with Sabakem Pty Ltd for the latest information on MRLs and import tolerances BEFORE using Sabakem Glufosinate 400 Herbicide

Resistance Warning:

GROUP 10 HERBICIDE

Sabakem Glufosinate 400 Herbicide is a member of the glycine group of herbicides. Sabakem Glufosinate 400 Herbicide has the inhibitor of glutamine synthetase mode of action. For weed resistance management Sabakem Glufosinate 400 Herbicide is a Group 10 herbicide. Some naturally occurring weed biotypes resistant to Sabakem Glufosinate 400 Herbicide, and other Group 10 herbicides which inhibit glutamine synthetase, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Sabakem Glufosinate 400 Herbicide or other Group 10 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Sabakem Pty Ltd accepts no liability for any losses that may result from the failure of Sabakem Glufosinate 400 Herbicide to control resistant weeds.

Precautions:

Precaution: DO NOT allow bystanders to come into contact with the spray cloud.

Re-entry period; Do not allow entry into treated areas until the spray has dried.

Protections:

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with this product or the used container.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT apply on desirable foliage or allow spray to drift onto the foliage of desirable plants, trees or vines, as damage will occur.

DO NOT allow product to contact green or uncalloused bark (such as on desirable young trees and vines) or cut, cracked, damaged or wounded tissue, where the affected surface is not adequately healed. Sabakem lufosinate 400 Herbicide may be used around desirable trees/vines less than two years old provided they are effectively shielded from spray and spray drift.

DO NOT allow desirable plant foliage to contact any inert surface, such as plastic mulches, which have been treated with Sabakem Glufosinate 400 Herbicide.

DO NOT apply Sabakem Glufosinate 400 Herbicide to recently fumigated or sterilised soil.

Storage and Disposal:

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. Do not re-use empty container for any other purpose.

Safety Directions:

Harmful if inhaled. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. Do not inhale spray mist.

When opening the container and preparing the spray, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves and goggles. If applying by boomspray equipment, wear cotton overalls, over normal clothing, buttoned to the neck and wrist, and elbow-length chemical resistant gloves. In addition, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, and a half piece respirator if applying by low pressure hand wand.

If product in eyes, wash it out immediately with water. Wash hands after use. After each days use wash gloves and goggles and contaminated clothing and respirator and if rubber wash with detergent and warm water.

First Aid Instructions:

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.

	A . I	1 A /	
Lirct	Λ 1Δ	Warn	inac:
	-	vvali	ings:

DIRECTIONS FOR USE

Crop / Situation	Weed	State	Rate	WHP	Critical Comments
Blackberry, boysenberry, loganberry, raspberry	Primocane and sucker control	NSW, Vic, Tas only	250mL/ 500L water	Nil	Apply as a directed spray to suckers and primocanes. Contact with flowers, developing fruit or desirable foliage will cause damage. Ensure complete coverage of primocanes/suckers by spraying to the point of runoff, preferably when they are less than 15cm high. Wetting agent may be added at a rate of 25mL/100L or equivalent.
Avocado, banana, feijoa, guava, kiwifruit, litchi, mango, pawpaw, passionfruit, pineapple, rambutan plantations	See list of weeds controlled in tables 1 and 2.	Qld, NSW, Vic, SA, WA, NT only	0.5 to 2.5 L/ha	Nil	Apply as a directed or shielded spray. Refer to the label section Application Equipment for specific information on application methods. Controlled Droplet Application equipment must not be used for application in cherry orchards. Warnings: DO NOT apply spray or spray drift to contact desirable foliage or green (uncalloused) bark. To avoid potential crop damage, refer to the label sections on Application Equipment and PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS. Sabakem Glufosinate 400 Herbicide may be used around trees/vines less than two years old provided they are effectively shielded from
Citrus orchards Olive plantations		All States			spray and spray drift. The recommended rate of use is determined by the following criteria: WEED SPECIES WEED GROWTH STAGE WEED DENSITY
Pome and stone fruit orchards Tree nut plantations Vineyards				21 days (H) Nil	CLIMATIC CONDITIONS WEED SPECIES Apply the appropriate rate to control the least susceptible weed present as per the lists of weeds controlled in the accompanying tables. WEED STAGE OF GROWTH Use the lower rate when weeds are young and succulent (grasses: pre-tillering; broadleaves: cotyledons to 4-leaf) or the population is very sparse. A median rate should be used for medium sized plants (grasses: tillering; broadleaves: 4-leaf to advanced vegetative) and the high rate should be used when weeds are mature (grasses: noding to flowering; broadleaves: budding to flowering).

Crop / Situation	Weed	State	Rate	WHP	Critical Comments
				-	WEED DENSITY
					Use the higher rates when the weed population is dense. Thorough coverage of weeds is essential for good control.
					CLIMATIC CONDITIONS
					Best results are achieved when applied under warm humid conditions. Control will be reduced and/or slower under cold conditions and/or overcast conditions. Good results will be achieved under most other conditions, however poor results may occur under hot dry conditions (temperature above 33°C with a relative humidity below 50%).
					Weeds that have been hardened or stunted in growth due to stressed conditions should be treated at the maximum rate.
					COVERAGE
					Complete coverage of weeds is essential for good control. Poor coverage may result in re-growth.
					PERENNIAL WEEDS Apply when weeds are actively growing. Follow-up treatments will be necessary to control re-growth of perennial weeds in most cases.
Strawberries, cane berry fruits (inter-row	See lists of weeds controlled	All states	0.5 to 2.5 L/ha	Nil	Apply as a directed or shielded spray to the inter-row area. Take care not to allow spray or spray drift to contact the crop, including strawberry runners. Refer
Tomatoes (inter-row)	in Tables 1 and 2				to GENERAL INSTRUCTIONS for warnings concerning plastic mulch and fumigated/sterilised soil. Determine the recommended rate of use by considering the criteria WEED SPECIES, WEED STAGE OF GROWTH, WEED DENSITY and CLIMATIC CONDITIONS, as described above.
Commercial & Industrial areas, rights-of-way and other non-agricultural areas including public service areas, roadsides, railways and beneath electrical transmission towers.	See lists of weeds controlled in Tables 1 and 2	All states	0.5 to 3.0 L/ha	-	Determine the recommended rate of use by considering the criteria WEED SPECIES, WEED STAGE OF GROWTH, WEED DENSITY and CLIMATIC CONDITIONS as described above. Warnings: Do not allow spray or spray drift to contact desirable plants. To avoid potential crop damage, refer to the label sections on Application Equipment and PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.

Optical Spot Spray Technologies (OSST)

Note: Calibrate the sprayer to spray the equivalent of 100 L/ha.

DO NOT apply greater than 30 L of spray mixture per hectare through OSST equipment (1.5 L of product per hectare).

SITUATION & CROP	WEEDS	WEED GROWTH STAGE	RATE	CRITICAL COMMENTS
Maintenance of summer fallow prior to planting:	Awnless barnyard grass (Echinochloa colona)	Up to 30 cm	5 L/100 L	Apply to actively growing weeds. Good coverage is essential. Refer to Application section for
Cereal grains (including wheat,	Bladder ketmia (Hibiscus trionum)	Up to 60 cm		details.
barley, oats, maize and sorghum);	Liverseed grass (Urochloa panicoides)	Up to 110 cm		DO NOT apply more than three applications per season.
Pulses (including chickpeas, faba beans, field peas,	Yellowvine (Tribulus micrococcus)	Up to 60 cm		DO NOT sow crops until 14 days or more have elapsed after the final application.
lentils, lupins and mung beans);	pigweed (Portulaca oleracea)	Up to 65 cm		Sabakem Glufosinate 400 Herbicide will affect weeds
Oilseeds (including canola, cotton, soybeans and sunflowers);				that are larger than the recommended leaf stage, but speed of activity and level of control may be reduced.
				CLIMATIC CONDITIONS Best results are achieved when Sabakem Glufosinate 400 Herbicide is applied under warm humid
				conditions (temperatures below 33°C with a relative humidity above 50%).
				Under any other conditions efficacy and speed of action may be reduced.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

Table 1. Recommendations for weed control (except when referred to Table 2).

		Application Rates			
Common Name	Scientific Name	Boom or Directed Sprayer L/ha	Handgun mL/100L	Knapsack mL/15L	
ANNUAL WE	1				
Amaranthus spp.	Amaranthus spp.	1.0 to 2.5	250	37.5	
Apple of Peru	Nicandra physalodes	0.75 to 1.5	150	22.5	
Argentine peppercress	Lepidium bonariense	1.0 to 1.5	150	22.5	
Awnless barnyard grass	Echinochloa colona	1.25 to 1.75	175	26.5	
Barley grass	Hordeum leporinum	1.0 to 1.5	150	22.5	
Barnyard grass	Echinochloa crus-galli	1.0 to 2.5	250	37.5	
Billy goat weed	Ageratum conyzoides	1.0 to 2.5	250	37.5	
Bitter cress	Cardamine hirsute	1.0 to 2.5	250	37.5	
Black bindweed (buckwheat) (refer Note 2)	Fallopia convolvulus	0.9 to 2.5	250	37.5	
Bladder ketmia	Hibiscus trionum	1.5 to 2.5	250	37.5	
Bordered panic	Entolasia marginata	1.0 to 2.0	200	30	
Brome grass (refer Note1)	Bromus spp.	1.0 to 1.5	150	22.5	
Calopo	Calopogonium mucanoides	1.0 to 2.5	250	37.5	
Caltrop burr (refer also Table 2)	Tribulus terrestris	1.5 to 2.5	250	37.5	
Capeweed	Arctotheca calendula	0.75 to 2.5	250	37.5	
Clover (subterranean)	Trifolium subterranean	0.9 to 1.5	150	22.5	
Cobbler's peg	Bidens pilosa	1.0 to 2.5	250	37.5	
Common storksbill	Erodium cicutarium	0.75 to 2.0	200	30	
Crowsfoot grass	Eleusine indica	1.5 to 2.5	250	37.5	
Deadnettle (refer also Table 2)	Lamium amplexicaule	1.0 to 2.5	250	37.5	
Dwarf crumbweed	Chenopodium pumilo	1.5 to 2.5	250	37.5	
Fat hen	Chenopodium album	1.5 to 2.5	250	37.5	
Fumitory	Fumaria officinalis	0.9 to 2.5	250	37.5	
Green crumbweed	Chenopodium carinatum	1.0 to 2.5	250	37.5	
Lesser canary grass (refer also Table 2)	Phalaris minor	1.5 to 2.5	250	37.5	
Liverseed grass (refer also Table 2)	Urochloa panicoides	0.75 to 2.5	250	37.5	
Medics (annual)	Medicago spp.	0.5 to 2.5	250	37.5	
Milk thistle	Sonchus oleraceus	1.0 to 2.5	250	37.5	
Mint weed	Salvia reflexa	1.5 to 2.5	250	37.5	
New Zealand spinach	Tetragonia tetragoniodes	1.0 to 2.5	250	37.5	
Patterson's Curse	Echium plantagineum	0.5 to 1.5	150	22.5	
Peanuts	Arachis hypogaea	0.75 to 1.5	150	22.5	
Pigweed	Portulaca oleracea	1.5 to 2.5	250	37.5	
Pinkburr	Urena lobata	1.0 to 2.5	250	37.5	
Potato weed	Galinsoga parviflora	1.0 to 2.5	250	37.5	
Praire grass (refer Note 1)	Bromus unioloides	2.0 to 2.5	250	37.5	
Prickly lettuce	Lactuca serriola	1.5 to 2.5	250	37.5	
Red natal grass	Rhynchelytrum repens	1.0 to 2.5	250	37.5	
Ryegrass (annual)	Lolium rigidum	1.0 to 2.5	250	37.5	
Saffron thistle	Carthamus lanatus	0.75 to 2.5	250	37.5	
St. Barnby's thistle	Centaurea solstitialis	0.75 to 2.5	250	37.5	
Sago weed	Plantago cunninghamii	1.0 to 1.5	150	22.5	
Scarlet pimpernel	Anagallis arvensis	1.0 to 2.5	250	37.5	
Setaria	Setaria italica	1.0 to 2.5	250	37.5	
Sheep thistle	Carduus tenuiflorus	1.25 to 2.5	250	37.5	
Silver grass	Vulpia myuros	1.0 to 2.5	250	37.5	
Sorghum/sudax	Sorghum bicolor	1.0 to 2.5	250	37.5	
Square weed	Spermacoce latifolia	1.0 to 2.5	250	37.5	
Stagger weed	Stachys arvensis	1.0 to 2.5	250	37.5	
Star of Bethlehem	Ipomoea quamoclit	1.0 to 2.5	250	37.5	
Summer grass	Digitaria cillaris	1.0 to 2.5	250	37.5	
Thickhead	Crassocephalum crepidioides	1.5 to 2.5	250	37.5	
Three Cornered Jack	Emex australis	1.0 to 2.5	250	37.5	

		Application Rates			
Common Name	Scientific Name	Boom or Directed Sprayer	Handgun mL/100L	Knapsack mL/15L	
Tomato	Lycopersicon esculentum	L/ha 1.0 to 2.5	250	37.5	
Turnip weed	Rapistrum rugosum	1.5 to 2.5	250	37.5 37.5	
Variegated thistle (refer also Table 2)	Silybum marianum	1.25 to 2.5	250	37.5 37.5	
Wheat	Triticum eastivum	2.0 to 2.5	250	37.5	
Wild carrot	Daucus glochidiatus	1.0 to 2.5	250	37.5	
Wild gooseberry	Physalis minima	1.0 to 2.5	250	37.5	
Wild mustrad	Sysimbrium orientale	1.0 to 2.5	250	37.5	
Wild oats (refer also Table 2)	Avena spp.	1.5 to 2.5	250	37.5	
Wild radish	Raphanus raphanistrum	2.5	250	37.5	
Wire weed (refer also Table 2)	Polygonum aviculare	0.75 to 2.5	250	37.5	
PERENNIAL V				,	
Blady grass	Imperata cylindrica	1.5 to 2.0	200	30	
Cape tulip	Homeria spp.	1.0 to 1.5	150	22.5	
Centro	Centrosema pubescens	0.5 to 2.5	250	37.5	
Clover glycine	Glycine latrobeana	0.5 to 1.5	150	22.5	
Couch grass	Cynodon dactylon	1.25 to 2.5	250	37.5	
Cow pea	Vigna unguiculata	0.5 to 1.5	150	22.5	
Giant sensitive plant	Mimosa invisa	1.0 to 2.5	250	37.5	
Greenleaf desmodium	Desmodium intortum	0.5 to 1.5	150	22.5	
Johnson grass	Sorghum halepense	1.5 to 2.5	250	37.5	
Panicum spp.	Panicum spp.	1.0 to 2.5	250	37.5	
Paspalum spp.	Paspalum spp.	1.5 to 2.5	250	37.5	
Perennial bindweed	Convolvulus arvensis	1.0 to 1.5	150	22.5	
Shamrock	Oxalis corymbosa	1.5	150	22.5	
Sida weed (refer also Table 2)	Sida retusa	1.5 to 2.5	250	37.5	
Silver leaf desmodium	Desmodium uncinatum	2.0 to 2.5	250	37.5	
Siratro	Macroptilium atropurpureum	0.5 to 1.5	150	22.5	
Stink grass	Eragrostis cilianensis	1.5 to 2.5	250	37.5	
White clover	Trifolium repens	1.5 to 2.5	250	37.5	
White eye	Richardia brasiliensis	1.5 to 2.5	250	37.5	
Willow herb	Epilobium spp.	2.0 to 2.5	250	37.5	

Notes: 1.

- 1. Well-established clumps of Prairie grass and Brome grasses may only be suppressed at these rates. Follow-up treatments may be necessary to control re-growth.
- 2. Good control will be achieved on small and medium sized plants only in non-crop situation.

Table 2. Commercial & Industrial areas, rights-of-way and other non- agricultural areas including public service areas, roadsides, railways and beneath electrical transmission towers (when referred from Table 1)

		A	Application Rate			
Common Name	Scientific Name	Boom or Directed Sprayer L/ha	Handgun mL/100L	Knapsack mL/15L		
	ANNUAL WEEDS					
Caltrop burr	Tribulus terrestris	2.0 to 2.5	250	37.5		
Dead nettle	Lamium amplexicaule	3.0	300	45		
Lesser canary grass	Phalaris minor	2.0 to 3.0	300	45		
Liverseed grass	Urochloa panicoides	0.75	75	11.5		
Variegated thistle	Silybum marianum	3.0	300	45		
Wild oats	Avena spp.	2.5 to 3.0	300	45		
Wire weed	Polygonum aviculare	1.0 to 2.5	250	37.5		
	PERENNIAL WEEDS					
Sida weed	Sida retusa	2.0 to 2.5	250	37.5		

GENERAL INSTRUCTIONS

Sabakem Glufosinate 400 Herbicide is a non-volatile herbicide with activity against many annual and perennial broadleaf weeds and grasses.

Sabakem Glufosinate 400 Herbicide is absorbed by plant foliage and green stems. It is not significantly translocated as an active herbicide throughout the plant, and therefore will only kill that part of a green plant that is contacted by spray. Sabakem Glufosinate 400 Herbicide does not provide residual weed control. Visible symptoms of control appear in 3 to 7 days, but complete desiccation may take 20 to 30 days under cool conditions.

Best results are achieved when application is made under good growing conditions.

Application to weeds under stress (e.g. due to continuous severe frosts, dry or waterlogged conditions) should be avoided.

Soil Fumigation / Sterilisation

Sabakem Glufosinate 400 Herbicide is metabolised (broken down) by microorganisms in the soil to become inactive. Soil fumigation or sterilisation will reduce the number of microorganisms present, thus slowing the breakdown of Sabakem Glufosinate 400 Herbicide.

As damage to transplants or seedlings may occur, it is not advisable to apply Sabakem Glufosinate 400 Herbicide in conjunction with soil fumigation or sterilisation.

Plastic Mulches

Sabakem Glufosinate 400 Herbicide will remain active on inert surfaces such as plastic. Special care should be taken when applying Sabakem Glufosinate 400 Herbicide over plastic mulches, as plant contact with the mulch after spraying may result in crop damage.

Compatibility

Sabakem Glufosinate 400 Herbicide is compatible with most residual herbicides e.g. simazine, diuron, oxyfluorfen, norfluazuron, and oryzalin, and with glyphosate and metsulfuron. The addition of a wetting agent or other adjuvant is generally not considered necessary, (refer to the Directions for Use table). However, benefit has been obtained using a wetting agent or adjuvant on hard-to-wet weeds when using water rates in excess of 500 L/ha. The rate is 25 mL/100 L of a 1000 g/L non-ionic wetting agent, or equivalent.

For information on compatible wetting agents and adjuvants, contact your local Sabakem Pty Ltd representative.

Mixing

Sabakem Glufosinate 400 Herbicide mixes easily with water. Clean water should always be used for mixing with Sabakem Glufosinate 400 Herbicide. Ensure that the spray tank is free of any residues of previous spray materials. Two-thirds fill the spray tank with clean water, and with agitator operating add the required amount of Sabakem Glufosinate 400 Herbicide.

Add other relevant compatible products. Top the tank up to the required volume with clean water with agitator running.

Application Equipment Ground Sprayers

Aim to apply a thorough and even coverage of spray to the target plant. Dense stands of weeds should be thoroughly wetted with spray. Incomplete coverage may result in poor control. Equipment should be such that adequate coverage, penetration and volume of spray liquid can be achieved.

Boom or Directed Sprayer Equipment

Sabakem Glufosinate 400 Herbicide should be applied at label rates (refer to specific column in the lists of weeds controlled) in sufficient water to give thorough coverage of weeds. It has been found that 300 to 500 L/ha has given good results under most weed conditions.

Special care must be taken when using sprayer/slasher combination units not to cause dust and turbulence, which can carry spray into non-target areas.

Knapsack and Handgun Equipment

Sabakem Glufosinate 400 Herbicide should be applied at label rates (refer to specific columns in the lists of weeds controlled) in adequate water to thoroughly wet the weeds being sprayed, i.e. 500 to 1000 L/ha. Dense stands will require up to 1000 L/ha of spray mixture, whereas less dense stands will

require less water. High volume application using hollow-cone nozzles for hand spraying is recommended.

Controlled Droplet Application (CDA) Equipment

Sabakem Glufosinate 400 Herbicide may be applied through CDA row spraying equipment fitted with a solid (impermeable) shroud or skirt, at rates as recommended for boom or directed sprayers (refer to specific column in the lists of weeds controlled), provided thorough spray coverage of weeds can be achieved. Apply preferably when weeds are less than 15 cm in height, with the equipment set up so that the spray dome only just touches the tops of the weeds. A total spray volume of 20 to 30 L/ha has been found to give good results. Do not mix residual herbicides or any spray adjuvants with Sabakem Glufosinate 400 Herbicide when using CDA equipment.

Warning: Because the spray solution is highly concentrated particular care must be taken when using Sabakem Glufosinate 400 Herbicide through CDA equipment to avoid contact of the spray solution with any part of the crop trunk or canopy. DO NOT apply Sabakem Glufosinate 400 Herbicide through equipment fitted with bristle skirts. Particular care should be taken when using CDA equipment around green or un-calloused bark.

Please refer to PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS. CDA equipment must not be used for application in cherry orchards.

Sprayer Cleanup

Clean all equipment after use by thoroughly flushing with water.

Aircraft

Do not apply by aircraft.