Product Name: Sabakem Buffer 700 APVMA Approval No: 94542/142729



Label Name:	Sabakem Buffer 700	
Signal Headings:	CAUTION	
	KEEP OUT OF REACH OF CHILDREN	
	READ SAFETY DIRECTIONS BEFORE OPENING OR USING	
Constituent	ACTIVE CONCILLIENTS	
Statements:	ACTIVE CONSTITUENTS	
	350 g/L SOYAL PHOSPHOLIPIDS 350 g/L PROPIONIC ACID	
Mode of Action:		
Statement of Claims:	Acidifying and Penetrating Surfactant Reduces Alkaline Hydrolysis of Dimethoate	
	Assists with Uptake of Foliar Fertilisers	
	Assists in Management of Spray Droplet Size	
Net Contents:	1L - 1000L	
Restraints:	DO NOT use with copper products.	
Directions for Use:	This section contains file attachment.	
Other Limitations:		

Withholding Periods:	
Trade Advice:	
General Instructions:	Sabakem Buffer 700 can be used where the addition of a wetter/spreader or self emulsifying oil is required or recommended on the agricultural chemical label.  Penetration and Wetter Sabakem Buffer 700 is a multipurpose adjuvant which is composed of natural surfactants and penetrants derived from soybean oils. These are combined with propionic acid to produce a penetrant, surfactant, acidifier for use with herbicides, insecticides, fungicides, foliar fertilizers and plant growth regulators. The unique properties of Sabakem Buffer 700 enhance the uptake of many systemic herbicides, allowing them to effectively penetrate into leaves, without causing damage to non-target crop plants.  Acidification Sabakem Buffer 700 also acts as an acidifier, which will reduce pH, in most cases to between 4 and 5. This reduces losses due to alkaline hydrolysis and also assists with the uptake of weak acid herbicides.  Droplet size management Sabakem Buffer 700 will reduce the number of fine droplets (<150micron) produced by flat fan nozzles, without increasing the number of large spray droplets (>400micron). Sabakem Buffer 700 will reduce the fine droplets associated with, but not eliminate, off-target movement of the pesticide being used. This is contingent upon good agricultural spraying practice and appropriate nozzle choice. DO NOT use in situations that are conducive to drift. Refer to Sabakem Pty Ltd for further advice and stewardship.  MIXING Half fill the spray tank with water and commence agitation. Add the required quantity of Sabakem Buffer 700 then add the recommended quantity of herbicide, insecticide or foliar fertiliser. Continue agitation while topping up the tank and during spraying. Do not use with copper products.
Resistance Warning:	
Precautions:	
Protections:	PROTECTION OF WILDLIFE, FISH CRUSTACEA AND ENVIRONMENT
	DANGEROUS TO FISH. DO NOT contaminate streams, rivers or watercourses with the chemical or used container.

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Storage and Disposal:

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemical on-site. Break, crush, or puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

## Safety Directions:

Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. When using together with other products, consult their label safety directions. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and face shield. Wash hands after use. After each day's use, wash gloves, face shield and contaminated clothing.

## First Aid Instructions:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766. If swallowed, DO NOT induce vomiting. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

First Aid Warnings:	:			
---------------------	---	--	--	--

## **DIRECTIONS FOR USE**

## **RATES OF APPLICATION**

Situation	Rate per 100L	Critical Comments
Addition to herbicides to improve spreading and penetration.	250 – 500 mL	Use high rate on stressed or difficult to control weeds.
Glyphosate, 2,4-D, MCPA, Metsulfuron-methyl		
Reduction of pH to reduce alkaline hydrolysis.	100 mL	Add to water in spray tank before adding Dimethoate.
Dimethoate		
To improve uptake of foliar fertilizers.	300 – 500 mL	Tank mixing with other agricultural chemicals may increase the potential for crop damage – check with supplier.
Assistance in droplet size management to partially reduce the number of fine droplets produced from hydraulic nozzles by air and ground.	300 – 500 mL	Helps to reduce the number of fine droplets. Sabakem Buffer 700 will reduce the fine droplets associated with, but not eliminate, off-target movement. This is contingent upon good agricultural spraying practice and appropriate nozzle choice. Refer to Sabakem representative for further advice and stewardship.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.