

Product Name: Sabakem Flumetsulam 800 WG Herbicide
APVMA Approval No: 94669/143145



Label Name:	Sabakem Flumetsulam 800 WG Herbicide
-------------	--------------------------------------

Signal Headings:	READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	ACTIVE CONSTITUENT: 800 g/kg FLUMETSULAM
-------------------------	--

Mode of Action:	<table><tr><td>GROUP</td><td>2</td><td>HERBICIDE</td></tr></table>	GROUP	2	HERBICIDE
GROUP	2	HERBICIDE		

Statement of Claims:	A water dispersible granule formulation for the post-emergence and salvage control of certain broadleaf weeds in winter cereals (including those undersown with clover, lucerne or medics); clover, fenugreek, lathyrus, lucerne, medic, serradella, and vetch (Popany only) seed crops and pastures; chickpeas, field peas, lentils, maize, peanuts; and for the pre-emergence control of certain broadleaf weeds in maize and soybeans as specified in the Directions For Use.
----------------------	--

Net Contents:	500 g - 15 kg
---------------	---------------

Restraints:	<p>DO NOT apply to plants which may be stressed (not actively growing) due to prolonged periods of extreme temperature (less than 5°C or greater than 30°C), moisture stress (water-logged or drought affected), poor nutrition or previous herbicide treatment as reduced levels of control may result.</p> <p>DO NOT apply post-emergence treatments if rain is likely within 4 hours.</p> <p>DO NOT irrigate (any method) treated crop or pasture for 48 hours after application.</p> <p>DO NOT apply to crops affected by disease or by previous herbicide treatment (eg triazines or sulfonylureas).</p> <p>Spray drift restraints</p> <p>Spray drift restraints Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.</p> <p>DO NOT allow bystanders to come into contact with the spray cloud.</p> <p>DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever</p>
-------------	---

	<p>possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.</p> <p>DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.</p> <p>DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings 1 to 2 hours before sunset and persist until 1 to 2 hours after sunrise.</p>
--	---

Directions for Use:	This section contains file attachment.
---------------------	--

Other Limitations:	
--------------------	--

Withholding Periods:	<p>WITHHOLDING PERIODS:</p> <p>HARVESTING WITHHOLDING PERIODS Chickpeas, field peas, lentils, maize, peanuts and soybeans: NOT REQUIRED WHEN USED AS DIRECTED Winter cereals: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION</p> <p>GRAZING/STOCK FOOD WITHHOLDING PERIODS Chickpeas, field peas, lentils, peanuts, soybeans, Popany vetch: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION. Barley, cereal rye, oats, triticale, wheat, grass pastures: DO NOT GRAZE FOR 3 DAYS AFTER APPLICATION. DO NOT CUT FOR STOCK FOOD OR HARVEST FOR SEED FOR 4 WEEKS AFTER APPLICATION. Maize: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION Clover, fenugreek, lathyrus, lucerne, medic, serradella: DO NOT GRAZE OR CUT FOR STOCK FOOD OR HARVEST FOR SEED FOR 3 DAYS AFTER APPLICATION</p>
----------------------	--

Trade Advice:	<p>EXPORT OF LIVESTOCK When SABAKEM FLUMETSULAM 800 WG HERBICIDE is used as directed and the above WHPs for grazing and cutting for stock food are observed, livestock fed treated commodities are considered acceptable to slaughter for export. However, export requirements are subject to change. Consult your exporter for updated information about specific export market requirements before feeding treated animal feeds to livestock.</p>
---------------	---

General Instructions:	This section contains file attachment.
-----------------------	--

Resistance Warning:	<p>RESISTANT WEEDS WARNING GROUP 2 HERBICIDE</p> <p>SABAKEM FLUMETSULAM 800 WG HERBICIDE is a broadleaf herbicide with no annual ryegrass activity and is a member of the triazolopyrimidine sulfonanilide (sulfonamide) group of herbicides. The product has the acetolactate synthase (ALS) inhibitor mode of action. For weed resistance management the product is a Group 2 herbicide. Some naturally occurring weed biotypes resistant to the product and other Group 2 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 2 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Sabakem Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Sabakem representative.</p>
---------------------	--

Precautions:	
--------------	--

Protections:	<p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS Refer to MINIMUM RECROPPING PERIODS for crop rotation information. Crops susceptible to SABAKEM FLUMETSULAM 800 WG HERBICIDE include but are not limited to canola, cotton, faba beans, lupins, sorghum and sunflowers. DO NOT flood irrigate any treated crop or pasture for 48 hours after application. Where other types of irrigation are used, for example sprinklers, DO NOT irrigate to the point of runoff for at least 48 hours after application. DO NOT apply to waterlogged soils or if heavy rain is expected within 48 hours of application. Dangerous to aquatic plants and susceptible crops. DO NOT contaminate dams, watercourses or drains with the product or its containers. DO NOT apply under weather conditions, such as dead calm or excessive wind, or from spraying equipment producing small droplets that may cause spray to drift onto adjacent areas, particularly wetlands, waterbodies, watercourses, susceptible crops or land to be planted with susceptible crops.</p> <p>PROTECTION OF LIVESTOCK DO NOT graze or cut treated crops for stock food except as specified under withholding periods. Poisonous plants may become more palatable after spraying, therefore livestock should be kept out of the area until the plants have died down.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.</p>
--------------	--

Storage and Disposal:	<p>KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a securely locked, dry, cool, well-ventilated area away from food, feedstuffs, fertilisers or seed. DO NOT store for prolonged periods in direct sunlight. DO Not re-use containers. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.</p>
-----------------------	--

	<p>If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p> <p>SMALL SPILL MANAGEMENT Sweep up material and contain in a refuse vessel for disposal in the same manner as for containers</p>
--	--

Safety Directions:	<ul style="list-style-type: none"> • Product will irritate the eyes. • When handling the granules avoid contact with eyes. • If product in eyes, wash it out immediately with water. • Wash hands after use.
--------------------	--

First Aid Instructions:	<p>If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.</p>
-------------------------	--

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE:

TABLE 1A. CHICKPEAS, FIELD PEAS, LENTILS, FENUGREEK, LATHYRUS, VETCH (POPANY ONLY) AND SERRADELLA

CROP	GROWTH	CROP TOLERANCE	SPRAY ADDITIVES/TANK
Chickpeas	4 - 6 branches (no later than 6 weeks after emergence)	Sabakem Flumetsulam 800 WG Herbicide usually causes some transient crop yellowing and can cause reddish discolouration and height suppression. Flowering may be delayed resulting in yield suppression.	DO NOT use any spray additives, or tank mix any other chemicals with Sabakem Flumetsulam 800 WG Herbicide when using on chickpeas and field peas.
Field Peas	2 to 6 nodes (no later than 6 weeks after emergence)	SABAKEM FLUMETSULAM 800 WG HERBICIDE may cause transient crop yellowing and height suppression. On light soils in dry seasons flowering may be delayed resulting in yield suppression.	
Lentils	4-8 fully expanded leaves DO NOT apply later than 6 weeks after crop emergence.	SABAKEM FLUMETSULAM 800 WG HERBICIDE may cause transient height reduction, crop discolouration and delayed flowering, although yields are normally unaffected. However, stress conditions after application (eg. frost, drought, nutrient deficiency, disease) may lengthen the time needed for lentils to recover. In seasons where a dry spring occurs, yields may be suppressed. Tank mixes with other products may result in growth suppression and delayed flowering which can result in yield suppression.	Uptake® Spraying Oil at 500 mL/100 L or BS-1000® at 200 mL/100 L may be applied with SABAKEM FLUMETSULAM 800 WG HERBICIDE to lentils.
Fenugreek, Lathyrus, Vetch (Popany only)	3 fully expanded leaves onwards		Use SABAKEM FLUMETSULAM 800 WG HERBICIDE or SABAKEM FLUMETSULAM 800 WG HERBICIDE plus a wetter only. Tank mixtures with other herbicides are not recommended.
Serradella	3 fully expanded leaves onwards		Uptake® Spraying Oil at 500 mL/100 L or BS-1000® at 200 mL/100 L may be applied with SABAKEM FLUMETSULAM 800 WG HERBICIDE for serradella.

TABLE 1B. WEEDS CONTROLLED OR SUPPRESSED IN TABLE 1A CROPS

WEED	WEED GROWTH STAGE		RATE g/ha	CRITICAL COMMENTS
	Up To Leaf No. or	Up To Plant size (cm)		
CONTROLLED				
Amsinckia (Yellow burrweed)	10 leaf	10 cm diameter	25	Where recommended, use of either a wetter or Uptake Spraying Oil with SABAKEM FLUMETSULAM 800 WG HERBICIDE will provide better weed control.
Ball mustard	6 leaf	5 cm diameter		
Charlock	8 leaf	10 cm diameter		Spray charlock as soon as possible after the autumn break. Larger plants and any affected by stress or grazing prior to treatment may re-grow and flower.
Indian hedge mustard	6 leaf	5 cm diameter		
Lupins	10 leaf	10 cm high		
Marshmallow (Small flowered mallow)	4 leaf	10 cm diameter		
Pheasant's eye	8 leaf	10 cm diameter		
Shepherd's purse	8 leaf	10 cm diameter		
Three-horned bedstraw	6 whorls	10 cm high		
Turnip weed	8 leaf	5 cm diameter		
Volunteer canola	8 leaf	10 cm diameter		
Ward's weed	8 leaf	10 cm diameter		
Wild turnip	6 leaf	5 cm diameter		
SUPPRESSED				
Capeweed (WA only)	4 leaf	10 cm diameter	25	Under ideal growing conditions SABAKEM FLUMETSULAM 800 WG HERBICIDE will provide useful suppression of capeweed and doublegee. Best results will be achieved when a pre- emergence herbicide has already been used.
Doublegee (Spiny emex) (WA only)	4 leaf	10 cm diameter		Under ideal growing conditions, SABAKEM FLUMETSULAM 800 WG HERBICIDE without an adjuvant will give a biomass reduction of 50% - 70% of wild radish. Surviving plants may flower and set viable seed. Best results will occur with treatment in conditions of >5°C with bright sunny conditions and use of higher water rates of 75-100 L/ha with fine-medium quality spray droplets to get excellent spray coverage.
Wild radish	4 leaf	5 cm diameter		

TABLE 2A. WHEAT, BARLEY, OATS, TRITICALE, CEREAL RYE (INCLUDING THOSE UNDERSOWN WITH CLOVER, LUCERNE OR MEDICS), CLOVER, LUCERNE AND MEDIC CROPS MIXED GRASSES/ LEGUME PASTURES

CROP / SITUATION	GROWTH STAGES	CROP TOLERANCE	SPRAY ADDITIVES/TANK MIXES
Wheat	3 leaf until start of jointing (Zadoks 13-31)		Always apply with Uptake Spraying Oil at 500 mL/100 L or a 100% concentrate non- ionic wetting agent such as BS-1000 at 200mL/100 L.
Barley, Oats	Mid-tillering to start of jointing (Zadoks 23-31)	Transient stem shortening and crop discolouration may occur, yields are normally unaffected. Where barley and oats are undersown, a vigorous legume component may lengthen the time needed for the cereal to recover, especially if the cereal is stressed by lack of moisture, trace element deficiency or disease. In severe cases, yields may be suppressed.	Use only with a wetting agent such as BS- 1000 when either applying SABAKEM FLUMETSULAM 800 WG HERBICIDE alone or with partner products in barley and oats.
Stirling barley (WA only)	Apply no earlier than Zadoks 31.		
Triticale, Cereal rye	Mid-tillering to start of jointing (Zadoks 23-31)		Always apply with Uptake Spraying Oil at 500 mL/100 L or a 100% concentrate non- ionic wetting agent such as BS-1000 at 200 mL/100 L.
Medic, lucerne, and clover seed crops and pastures, including; Barrel medic, Snail medic, Spineless burr medic, Subterranean clover, White clover	2 to 3 trifoliolate leaves onwards	Medic, lucerne and subterranean clover (sub clover) - When SABAKEM FLUMETSULAM 800 WG HERBICIDE is applied at 25 g/ ha + Uptake or wetter, yield reduction may occur when treating Serena medic or Nungarin sub clover. DO NOT apply to lucerne seed crops less than 8 weeks before flowering	Always apply with Uptake Spraying Oil at 500 mL/100 L or a 100% concentrate non- ionic wetting agent such as BS-1000 at 200 mL/100 L. In lucerne DO NOT use with MCPA. In medics DO NOT use with MCPA.
Grazing lucerne - high rate	4 trifoliolate leaves onwards	Use the 50 g/ha rate in grazing lucerne only. DO NOT apply at 50 g/ha to lucerne used for seed production.	
Fence lines, Stock camps, Stockyards, Commercial areas and pastures including medic, lucerne and clover pastures.	2 to 3 trifoliolate leaves onwards (see crop tolerance)		Use Uptake Spraying Oil at 500 mL/100 L.

SALVAGE SPRAY			
Cereals: Wheat Barley Oats, Triticale, Cereal rye	Flowering (anthesis) to early dough (Zadoks, 61-83)		
Pastures, Lucerne, Clover, Medics	Advanced seedlings or re-growth after cutting or grazing		

TABLE 2B. WEEDS CONTROLLED IN TABLE 2A CROPS

WEED	WEED GROWTH STAGE		RATE g/ha	CRITICAL COMMENTS
	Up To Leaf No. or	Up To Plant size (cm)		
Amsinckia (Yellow burrweed)	10 leaf	10 cm diameter	25 + Uptake or wetter	
Ball mustard	6 leaf	5 cm diameter		
Buchan weed	8 leaf	10 cm diameter	Lucerne and/or clover only 25 + wetter + 2,4-DB (500 g/L) 1.5-2.5 L/ha	
			Grazing lucerne only 50 + Uptake or wetter	Use the 50 g/ha rate in grazing lucerne only and apply from 4 trifoliate leaves onwards. DO NOT apply at 50 g/ha to lucerne used for seed production.
Calepina (White ball mustard)	8 leaf	10 cm diameter	25 + Uptake or wetter	
Capeweed	4 leaf	10 cm diameter	25 + Uptake or wetter + bromoxynil (200 g/L) 700 mL/ ha	Optimum results are obtained in a competitive pasture. For best results follow up with moderate grazing two weeks after application. In pasture, spray as soon as possible after the autumn break. Larger plants and any affected by stress or grazing prior to treatment may re-grow and flower.
Charlock	8 leaf	10 cm diameter	25 + Uptake or wetter	Spray as soon as possible after the autumn break. Larger plants and any affected by stress or grazing prior to treatment may re-grow and flower.
Cotula (WA only)	4 leaf	10 cm diameter	25 + Uptake or wetter	Use Uptake Spraying Oil at 500 mL/100 L or wetting agent such as BS-1000 at 200 mL/100 L.

Doublegee (Spiny emex)	6 leaf	15 cm diameter	25 + Uptake or wetter + bromoxynil (200 g/L) 700 mL/ ha or 25 + wetter + 2,4-DB (500 g/L) 1.5 - 2.5 L/ha	Optimum results are obtained in a competitive pasture. For best results follow up with moderate grazing two weeks after application. In pasture, spray as soon as possible after the autumn break. Larger plants and any affected by stress or grazing prior to treatment may re-grow and flower. Use Uptake Spraying Oil or a wetter with SABAKEM FLUMETSULAM 800 WG HERBICIDE + bromoxynil tank mixture.
Dwarf marigold (Poverty weed)	10 leaf	15 cm high	15 + Uptake or wetter	
Fat hen	15 leaf	20 cm high	Spring/summer pasture and lucerne only 25 + Uptake or wetter	Spring and summer pasture and lucerne application only.
			Grazing lucerne only 50 + Uptake or wetter	Use the 50 g/ha rate in grazing lucerne only and apply from 4 trifoliolate leaves onwards. DO NOT apply at 50 g/ha to lucerne used for seed production.
Fumitory	6 leaf	8 cm diameter	25 + 300 ml/ha terbutryn (500 g/L) + wetter	Only use a wetter with SABAKEM FLUMETSULAM 800 WG HERBICIDE + terbutryn tank mixes. Note: this mixture is only approved for use in NSW, Vic and Tasmania on pastures.
Hedge mustard	8 leaf	10 cm diameter	25 + Uptake	
Lupins	10 leaf	10 cm high	25 + Uptake or wetter	
WA blue and narrow leaf lupins	4 to 8 leaf		10 + Uptake or wetter	
Marshmallow (Small flowered mallow) seedlings	4 leaf	10 cm diameter	25 + Uptake or wetter or 15 + wetter + 700 mL/ha bromoxynil-MCPA (200 g/L + 200 g/L), or 15 + wetter + 350 mL/ha terbutryn (500 g/L) + 700 ml/ha MCPA amine (500 g/L)	Add a wetter to MCPA/terbutryn or bromoxynil-MCPA mixes. Only use bromoxynil/MCPA and terbutryn + MCPA mixes in cereals that are not undersown with clovers, medics or lucerne.
	10 leaf	20 cm diameter	25 + wetter + 2,4-DB (500 g/L) 1.5-2.5 L/ha	For older plants see Weeds Suppressed. only use a wetter with SABAKEM FLUMETSULAM 800 WG HERBICIDE + 2,4 D,B tank mixes.
			Grazing lucerne only 50 + Uptake or wetter	Use the 50 g/ha rate in grazing lucerne only and apply from 4 trifoliolate leaves onwards. Do not apply at 50 g/ha to lucerne used for seed production.

Paterson's curse (Salvation Jane)	8 leaf	10 cm diameter	25 + Uptake or wetter + bromoxynil (200 g/L) 700 ml/ha or 25 + wetter + terbutryn (500 g/L) 300 mL/ha	In pasture, larger plants and any affected by stress or grazing prior to treatment may re-grow and flower. For best results follow up with moderate grazing two weeks after application. With terbutryn, apply in a minimum spray volume of 100 L/ha from the ground or 50 L/ha from aircraft.
Peppercress seedlings	8 leaf	10 cm diameter	25 + Uptake or wetter	
	10 leaf	15 cm diameter	25 + wetter + 2,4-DB (500 g/L) 1.5 - 2.5 L/ha	
Pheasant's eye	7 leaf	10 cm high	25 + Uptake or wetter	
Shepherd's purse	8 leaf	10 cm diameter		
Three-horned bedstraw	6 whorls	10 cm high		
Turnip weed	8 leaf	5 cm diameter	15 + Uptake or wetter	
	12 leaf	10 cm diameter	25 + Uptake or wetter	
Volunteer canola	8 leaf	10 cm diameter		
Ward's weed				
Wild radish	6 leaf	15 cm diameter	25 + Uptake or wetter + bromoxynil (200 g/L) 700 mL/ha or 25 + wetter + MCPA amine (500 g/L) 500 mL/ha	When conditions at spraying are less than ideal (see RESTRAINTS above), or when the crop is not competitive, some radish plants may survive to flower and set viable seed.
Wild radish (cereals)	6 leaf	15 cm diameter	15 + wetter + 700 mL/ha bromoxynil/MCPA (200 g/L + 200 g/L) or 15 + Uptake or wetter + 700 mL/ha MCPA amine (500 g/L) or 15 + wetter + 700 mL/ha MCPA amine (500 g/L) + 350 mL/ha terbutryn (500 g/L)	DO NOT use MCPA amine in cereals undersown with clover, medics or lucerne. In lucerne DO NOT use MCPA. In medics DO NOT use MCPA Add Uptake Spraying Oil or wetter to SABAKEM FLUMETSULAM 800 WG HERBICIDE mixes with MCPA amine and a wetter to SABAKEM FLUMETSULAM 800 WG HERBICIDE + MCPA/terbutryn or SABAKEM FLUMETSULAM 800 WG HERBICIDE + bromoxynil/MCPA mixes. DO NOT use MCPA amine or MCPA amine + terbutryn in cereals undersown with clover, medics or lucerne.
Wild turnip	10 leaf	10 cm diameter	25 + Uptake or wetter	

Wireweed	10 leaf	15 cm diameter	Pasture and lucerne only 25 + wetter +2,4-DB (500 g/L) 1.5-2.5 L/ha	Undersown clovers and Lucerne, spring and summer sown pasture and lucerne crops only.
----------	---------	----------------	--	---

TABLE 2C. WEEDS SUPPRESSED IN TABLE 2A CROPS

WEED	WEED GROWTH		RATE g/ha	CRITICAL COMMENTS
	Up To Leaf No. or	Up To Plant size (cm)		
Buchan weed	8 leaf	10 cm diameter	25 + Uptake or wetter	Only use a wetter with SABAKEM FLUMETSULAM 800 WG HERBICIDE +2, 4-DB tank mixes.
Deadnettle	6 leaf	5 cm diameter	25 + wetter + 2,4-DB (500 g/L) 1.5 - 2.5 L/ha	
Doublegee (Spiny emex)	4 leaf	10 cm diameter	25 + Uptake or wetter	
	6 leaf	15 cm diameter	Grazing lucerne only 50 + Uptake or wetter	Use the 50 g/ha rate in grazing lucerne only and apply from 4 trifoliolate leaves onwards. DO NOT apply at 50 g/ha to lucerne intended for seed production.
Marshmallow (Small flowered mallow)	5-8 leaf	10 cm diameter	25 + Uptake or wetter	Only use a wetter with SABAKEM FLUMETSULAM 800 WG HERBICIDE + 2,4-DB tank mixes.
New Zealand spinach	4 leaf	5 cm diameter		
Paterson's curse (Salvation Jane)	8 leaf	10 cm diameter		
Peppercress	10 leaf	15 cm diameter	25 + Uptake or wetter	
Stagger weed	6 leaf	5 cm diameter	25 + wetter + 2,4-DB (500 g/L) 1.5 - 2.5 L/ha	
Wild radish	4 leaf	5 cm diameter	25 + Uptake or wetter	

TABLE 3A. DIURON TANK MIXES: WHEAT, BARLEY, OATS, TRITICALE, CEREAL RYE

Tank Mixtures: Read all label directions including restraints, spray drift restraints, mandatory no-spray zones, critical comments, withholding periods, regional use restrictions and safety Directions for the tank mix products.

CROP/SITUATION	GROWTH STAGES	CROP TOLERANCE	SPRAY ADDITIVES/TANK MIXED
Wheat	3 leaf until start of jointing (Zadoks 13-31)		Always apply with Uptake Spraying oil at 500 mL/100 L or a 100% concentrate non-ionic wetting agent such as BS-1000 at 200 mL/100L.
Barley Oats	Mid-tillering to start of Jointing (Zadoks 13-31)	Transient stem shortening and crop discolouration may occur, although yields are normally unaffected. In severe cases yields may be	Use only with a wetting agent such as BS-1000 when either applying SABAKEM FLUMETSULAM 800 WG
Stirling Barley (WA)	Apply no earlier than Zadoks 31		

		suppressed	HERBICIDE alone or with partner products in barley and oats.
Triticale Rye	Mid-tillering to start of Jointing (Zadoks 13-31)		Always apply with Uptake Spraying oil at 500 mL/100 L or a 100% concentrate non-ionic wetting agent such as BS-1000 at 200 mL/100L.
SALVAGE SPRAY			
Cereals: Wheat Barley Oats Triticale Cereal Rye	Flowering (anthesis) to early dough (Zadoks 61-83)	DO NOT use more than 25 g/ha	

TABEL 3B. DIURON TANK MIXES: WEEDS CONTROLLED IN TABLE 3A

WEED	WEED GROWTH STAGE		RATE g/ha	CRITICAL COMMENTS
	Up to leaf No. or	Up to plant size (cm)		
Capeweed	4 leaf	10 cm diameter	25 + wetter + 100 mL/ha diuron (500 g/L)	Optimum results are obtained in a competitive pasture. For best results follow up with moderate grazing two weeks after application. Larger plants and any affected by stress or grazing prior to treatment may re-grow and flower.
Doublegee (Spiney emaex)	4 leaf	10 cm diameter	25 + wetter + 100 mL/ha diuron (500 g/L)	

TABLE 4. AGRICULTURAL NON-CROP AREAS

WEED	WEED GROWTH STAGE	RATE g/ha	CRITICAL COMMENTS
Caltrop, Capeweed, Marshmallow (Small flowered mallow) (suppression), Paterson's curse (Salvation Jane), Wild radish	Rosette stage prior to running up to flower	Spot spray: 25 g/100 L	Apply to actively growing rosettes. To ensure complete coverage, spray to the point of runoff. Use Uptake Spraying Oil at 500 mL/100 L.

TABLE 5A. SEED CROPS (Tasmania only): SUBTERRANEAN CLOVER, RED CLOVER, WHITE CLOVER, ARROWLEAF CLOVER AND LUCERNE.

CROP	GROWTH STAGES	CROP TOLERANCE	SPRAY ADDITIVES/TANK MIXES
Seed crops of Subterranean clover, Red clover, White clover, Arrowleaf clover, Lucerne.	1 to 3 trifoliate leaves onwards	DO NOT apply to lucerne or clover seed crops less than 8 weeks before flowering. DO NOT apply at 40 g/ha to lucerne intended for seed production.	Use Uptake Spraying Oil at 500 mL/100 L or a wetting agent such as BS-1000 at 200 mL/100 L. In clover and lucerne SABAKEM FLUMETSULAM 800 WG HERBICIDE may be tank-mixed with 2,4-DB and/or bromoxynil at their respective label rates for complete control of suppressed weeds.

TABLE 5B. WEEDS CONTROLLED OR SUPPRESSED IN TABLE 5A CROPS

WEED	WEED GROWTH STAGE	RATE g/ha	CRITICAL COMMENTS
WEEDS CONTROLLED			
Charlock	Up to 3½ leaf stage	25 + Uptake or wetter	Use Uptake Spraying Oil at 500 mL/100 L or wetting agent such as BS-1000 at 200 mL/100 L.
Fat hen, Lesser swinecress, Mustards, Shepherd's purse, Wild radish, Wild turnip	Beyond 3½ leaf stage and up to 10 leaf stage	40 + Uptake or wetter	
WEEDS SUPPRESSED			
Capeweed Chickweed Fumitory Spurrey Wireweed	Beyond 3½ leaf stage and up to 10 leaf stage	40 + Uptake or wetter	<p>In clover and lucerne, seedlings of these weeds will be suppressed with SABAKEM FLUMETSULAM 800 WG HERBICIDE alone.</p> <p>In clover and lucerne, SABAKEM FLUMETSULAM 800 WG HERBICIDE may be tank-mixed with 2,4-DB and/or bromoxynil at their respective label rates for complete control of suppressed weeds. Only use a wetting agent at 200 mL/100 L with these tankmixes.</p>

TABLE 6A. SOYBEANS, LUCERNE, MAIZE AND PEANUTS

CROP	GROWTH STAGES	APPLICATION METHODS	SPRAY ADDITIVES / TANK MIXED	CROP TOLERANCE
Maize	Post-plant pre-emergence (PPPE)	Apply SABAKEM FLUMETSULAM 800 WG HERBICIDE after planting and before emergence of crop and weeds. Apply to moist soil only.	May be tank mixed with pendimethalin.	
Soybean	Pre-plant Incorporated (PPI)	Incorporate into the soil within 4 hours by making two passes in opposite directions using a combine with trailing harrows or similar equipment, to ensure thorough incorporation.	May be tank mixed with Trifluralin or pendimethalin.	Some transitory crop yellowing and height suppression should be expected but yields will be unaffected.
	Incorporated By Sowing (IBS)	Ensure the planting operation is done within 4 hours of application, using a combine with trailing harrows or similar equipment, to ensure thorough incorporation.	May be tank mixed with Trifluralin or pendimethalin.	
	Post-plant Pre-emergent (PPPE)	Apply SABAKEM FLUMETSULAM 800 WG HERBICIDE after planting and before emergence of crop and weeds. Apply to moist soil only.	May be tank mixed with pendimethalin.	
Lucerne	Post-emergent Up to 6 trifoliate leaf stage		DO NOT apply at 50 g/ha to lucerne intended for seed production. Apply with Uptake Spraying Oil at 500 mL/100 L spray volume or with a 100% concentrate non-ionic wetter such as BS-1000 at 200 mL/100 L.	
Peanuts	Post-emergent Up to 6 leaf stage			

TABLE 6B. WEEDS CONTROLLED OR SUPPRESSED IN TABLE 6A CROPS

WEED	WEED GROWTH STAGE	RATE g/ha	CRITICAL COMMENTS
WEEDS CONTROLLED			
Annual ragweed, Boggabri Weed, Fat hen, Wild radish (IBS and PPPE only)	Pre-emergent	25 or 50	WEED CONTROL: Minimum spray volume 150 L/ha for optimum results. In pre-emergent situations use the higher rate for longer soil residual effect and better suppression of more tolerant weeds (see WEEDS SUPPRESSED). In post-emergent situations use 25 g/ha on weeds up to 2 leaf stage and 50 g/ha on larger weeds up to 4 leaf stage and where more residual control is required.
Caltrop, Fat hen Turnip weed, Wild radish	Post-emergent Up to 4 leaf	25 or 50 + Uptake or wetter	

WEEDS SUPPRESSED			
Black pigweed, Bladder ketmia, Caltrop, Cobbler's pegs	Pre-emergent	25 or 50	WEED CONTROL: Minimum spray volume 150 L/ha for optimum results. In pre-emergent situations use the higher rate for longer soil residual effect and better suppression of more tolerant weeds (see WEEDS SUPPRESSED). In post-emergent situations use 25 g/ha on weeds up to 2 leaf stage and 50 g/ha on larger weeds up to 4 leaf stage and where more residual control is required.
Annual ground cherry, Anoda weed, Bladder Ketmia, Boggabri weed, Fierce thornapple (Qld only), Red pigweed, Wild gooseberry	Post-emergent Up to 4 leaf	25 or 50 + Uptake or wetter	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

GENERAL INSTRUCTIONS:

MINIMUM RECROPPING PERIODS

Cereal rye, medics, triticale, wheat, maize, soybeans	May be planted at any time after application of CropSure Counterstrike 800WG Herbicide
Barley, chickpeas, clover, field peas, Lucerne, oats and peanuts	Allow 3 months to elapse after application before sowing these crops
Canola, cotton, faba beans, fenugreek, lathyrus, lentils, lupins, serradella, sorghum, sunflowers, Popany vetch	On deep soils (with no impermeable sub-horizon), cotton, sorghum and sunflowers may be planted 3 months after application of CropSure Counterstrike 800WG Herbicide. Canola, faba beans and lupins are more sensitive and may be planted 9 months after application of CropSure Counterstrike 800WG Herbicide. On shallow, duplex, low organic matter soils with an impermeable sub-horizon within the root zone (30cm deep or less) these crops should NOT be planted until 2 years after application of CropSure Counterstrike 800WG Herbicide.

MIXING

Quarter fill the spray tank and add the required amount of Sabakem Flumetsulam 800 WG Herbicide. Add the remaining water with the agitator running. Add Uptake Spraying Oil or the wetting agent last (if used). Maintain agitation during spraying. Only mix sufficient spray solution for immediate use and avoid storing.

When tank mixing: SABAKEM FLUMETSULAM 800 WG HERBICIDE should be added to the tank first, followed by wettable powders or other dry flowable formulations, suspension concentrates (flowables), aqueous concentrates (eg. Lontrel* Herbicide), emulsifiable concentrates (eg. Verdict* 520 Herbicide) and then add Uptake Spraying Oil or wetting agent last (if used).

APPLICATION

Apply SABAKEM FLUMETSULAM 800 WG HERBICIDE in 50 to 150 litres of water per hectare, through an accurately calibrated boom sprayer. For aircraft application apply SABAKEM FLUMETSULAM 800 WG HERBICIDE in no less than 30 L/ha of water through accurately calibrated equipment. The product should be applied by an accurately calibrated ground rig or aircraft delivering medium quality spray. Best results are achieved where applications are made on warm (greater than 5°C), sunny days applying more than 50 L/ha of total spray volume (preferably more than 75 L/ha) and where spray coverage is maximised.

COMPATIBILITY

- Always allow 7 days between application of a grass herbicide and SABAKEM FLUMETSULAM 800 WG HERBICIDE in chickpeas and field peas.
- In lentils, adjuvant, broadleaf or grass herbicide, insecticide and foliar fertiliser tank mixes may result in transient height reduction, crop discolouration and delayed flowering, although yields are normally unaffected. However, stress conditions after application (eg. frost, drought) may lengthen the time needed for lentils to recover and in years where a dry spring occurs, yields may be suppressed.

CLEANING SPRAY EQUIPMENT

After using SABAKEM FLUMETSULAM 800 WG HERBICIDE, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose. Drain the tank and clean any filters in the tank, pump, lines and nozzles.

- **To rinse:** After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

- **To decontaminate:** Before spraying sensitive crops (which include canola, cotton, faba beans, lupins, sorghum and sunflowers), wash the tank and rinse the system as above. Quarter fill the tank and add an alkali detergent (e.g. SURF®, Cold Water SURF Concentrate®, Dynamo Matic Concentrate®, OMO® or DRIVE® at 500 mL/100 L of water or the powder equivalent at 500 g/100 L) and circulate throughout the system for at least fifteen minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain. Chlorine based cleaners are not recommended. Nufarm Tank and Equipment Cleaner® is not recommended.

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.